

Thermodynamics of diffusive DM/DE systems

Z. Haba

Institute of Theoretical Physics, University of Wrocław,
50-204 Wrocław, Plac Maxa Born'a 9, Poland
email:zhab@ift.uni.wroc.pl

December 20, 2016

Abstract

We discuss the energy density, temperature and entropy of dark matter (DM) and dark energy (DE) as functions of the scale factor a in an expanding universe. In a model of non-interacting dark components we repeat a derivation from thermodynamics of the well-known relations between the energy density, entropy and temperature. In particular, the entropy is constant as a consequence of the energy conservation. We consider a model of a DM/DE interaction where the DM energy density increase is proportional to the particle density. In such a model the dependence of the energy density and the temperature on the scale factor a is substantially modified. We discuss (as a realization of the model) DM which consists of relativistic particles diffusing in an environment of DE. The energy gained by the dark matter comes from a cosmological fluid with a negative pressure. We define the entropy and free energy of such a non-equilibrium system. We show that during the universe evolution the entropy of DM is increasing whereas the entropy of DE is decreasing. The total entropy can increase (in spite of the energy conservation) as the DM and DE temperatures are different. We discuss non-equilibrium thermodynamics on the basis of the notion of the relative entropy.

1 Introduction

The universe evolution is determined by Einstein equations with an energy-momentum tensor on their rhs. The energy-momentum is to contain an information about a large system of "particles" (galaxies, interstellar gas, dark matter etc.). A deterministic description of such a system is unrealistic. The successful Λ CDM model [1] determines the global evolution of the universe. In order to go into its more detailed structure we must construct particular models of the energy-momentum (cosmological fluids, inflaton fields). Simple models involve non-interacting components of the energy-momentum. However, if we

wish to explain a relation between the components and their evolution in time we need to take into account an interaction between components. Some basic properties of large systems are a consequence of their statistical and thermodynamic description (as their macroscopic version). The thermodynamic point of view has been developed from the beginning of cosmology [2]. It concerned non-interacting fluids. In this paper we discuss some consequences of an interaction. Einstein equations have the energy-momentum tensor on their rhs which is conserved. As a consequence, in a homogeneous universe the entropy of matter consisting of non-interacting components is also conserved. However, if the energy momentum consists of components which exchange energy then the corresponding entropies of the subsystems change in time. We consider a system consisting of radiation, "baryonic" matter, dark matter (DM) and dark energy (DE). The energy momentum of "baryonic" matter and radiation is conserved. The energy momentum of the dark matter is not conserved and its divergence has the opposite sign with respect of the divergence of the energy momentum of the dark energy. We propose a particular model of the energy-momentum non-conservation. The change of the energy-density is proportional to the particle density. In sec.3 we study some thermodynamic consequences of this assumption. In sec.4, based on our earlier papers [3][4] we introduce a particular realization of this energy-momentum non-conservation resulting from a diffusion of DM in a fluid of DE. In sec.2 we review some old derivations of the relations between energy density, temperature and entropy (the derivation follows the text-book of Planck [5]) as a consequence of the first law of thermodynamics or the energy-momentum conservation. In sec.3 we show how these relations are changed if we have components of the energy-momentum which are not conserved. We derive a formula for the energy density. On a basis of general thermodynamic principles we suggest a formula for a dependence of the temperature on the scale factor a of an expanding universe. Then, using the Clausius-Caratheodory concept of the temperature as the integrating factor for the heat we can calculate the entropy of each component of the cosmological fluid. The entropy of the universe has been discussed in many papers [2][6][7][8][9] [10][11][12]. The discussion concerned mainly non-interacting fluids. The large value of the estimated entropy of the universe poses a serious problem of its origin. It clearly contradicts the mechanical "adiabatic" evolution of the universe based on the Λ CDM model. In our model the total entropy of the fluids is increasing in spite of the energy conservation (in contradistinction to non-interacting fluids). In sec.5 we discuss thermodynamics of diffusing systems using the notion of relative entropy [13][14]. It allows to follow the relation between entropy, energy and the free energy. There are many computer simulations of structure formation in which the entropy of the components of the cosmic fluid gives an important information about the future evolution of the system and exchange of heat inside the system [15] [16]. In this paper, in a particular model, we study the temperature of the components and an evolution of their entropy. In sec.6 we discuss also the generalized second law of ther-

modynamics [17][18][19] (taking into account the entropy of the horizon). The entropy of the horizon is the dominating contribution to the universe entropy [11][12], but it does not change the conclusions about the entropy increase in the models of diffusive interactions. In the Appendix we show how to extend our results to an arbitrary DM/DE interaction.

2 Thermodynamics of non-interacting DM/DE systems

As an introduction, let us begin with a simplified description of equilibrium thermodynamics and its extension to an expanding universe (for a more general approach see [20]). Let us consider the first law of thermodynamics in the form

$$TdS = dE + pdV, \quad (1)$$

where S is the entropy, T temperature, E energy and p is the pressure. We define the energy density ρ by

$$E = \rho V, \quad (2)$$

where V is the volume. Let us assume the equation of state

$$p = w\rho, \quad (3)$$

where w is a constant. Then, treating $S(T, V)$ as a function of V and T we have

$$\left(\frac{\partial S}{\partial T}\right)_V = \frac{V}{T} \frac{d\rho}{dT}. \quad (4)$$

$$\left(\frac{\partial S}{\partial V}\right)_T = (1 + w) \frac{\rho}{T}. \quad (5)$$

Comparing second order partial derivatives we get

$$\left(\frac{\partial \rho}{\partial T}\right)_V = w^{-1}(1 + w) \frac{\rho}{T}. \quad (6)$$

Hence,

$$\rho(T) = BT^{\frac{1+w}{w}} \quad (7)$$

and from eq.(5)

$$S = (1 + w)BVT^{\frac{1}{w}}. \quad (8)$$

with a certain constant B . As a consequence, for a negative $w > -1$ the energy density ρ as well as the entropy S are decreasing functions of temperature. The derivation of eqs.(4)-(8) for $w = \frac{1}{3}$ appeared already in the text-book of Planck [5] (p.62). The formulas can still apply in an expanding universe. If we demand that in an expanding flat universe with $V = V_0 a^3$ (where a is the expansion

scale factor) the entropy is constant (an adiabatic expansion) then from eq.(8) ($a(0) = 1$)

$$T_a = T_0 a^{-3w}. \quad (9)$$

We obtain from eq.(9) the well-known a^{-1} decrease of the temperature for radiation (with $w = \frac{1}{3}$).

Let us show that eq.(9) is in agreement with the energy-momentum conservation

$$(T^{\mu\nu})_{;\mu} = 0 \quad (10)$$

and its representation as an ideal fluid

$$T^{\mu\nu} = (\rho + p)u^\mu u^\nu - pg^{\mu\nu} \quad (11)$$

with a four-velocity $u^\mu u_\mu = 1$. After choosing the comoving frame $u = (1, 0)$ the conservation of the energy-momentum tensor takes the form

$$\partial_t \rho + 3H(\rho + p) = 0, \quad (12)$$

where $H = a^{-1}\partial_t a$. Inserting $V = V_0 a^3$ in eq.(1) we derive from eq.(12)

$$dS = 0 \quad (13)$$

confirming that the entropy is a constant and as a consequence confirming also eq.(9). Moreover, we can solve eq.(12) with the result

$$\rho = \rho_0 a^{-3(1+w)}. \quad (14)$$

Comparing with eq.(7) we get again the formula (9) (which was derived from eq.(13) as resulting from the energy density expansion law). So, the temperature law (7) (Stefan-Boltzmann formula for $w = \frac{1}{3}$) follows either from the conservation of energy or from the conservation of entropy.

If we have a non-interacting DM/DE system of ideal fluids then the entropy of each component is constant. With $\tilde{w} > 0$ for DM and $w < 0$ for DE the density of DM is decreasing in time faster than the density of DE (the density of the phantom DE with $w < -1$ is increasing). The temperature of DM is decreasing whereas the temperature of DE is increasing. Hence, there cannot be any equilibrium between DM and DE. We have also the coincidence problem that the densities of DM and DE are of the same order only at the present time.

It is known that thermodynamics of non-ideal or inhomogeneous fluids substantially modifies the temperature and energy density evolutions [6][21][22][23]. We may expect that an interaction of fluids will also have a profound effect of the thermodynamics of components. This could be seen by a division of a non-homogeneous fluid into homogeneous interacting parts.

3 Thermodynamics of the DM/DE interaction

There are many models of DM/DE interaction (see [24] for a review of interacting fluids). The thermodynamics of a non-conserved energy-momentum in the cosmological models has been discussed in [25][26][27]. We consider in this section a particular class of models when the energy-momentum non-conservation is expressed by the particle density.

We restrict our discussion to the metric

$$ds^2 = g_{\mu\nu} dx^\mu dx^\nu = dt^2 - a^2 h_{jk} dx^j dx^k \quad (15)$$

(mainly to the FLWR metric when the tensor h_{jk} describes the homogenous space of constant curvature). The energy-momentum tensor $\tilde{T}$ of the dark matter under general assumptions can be expressed in the form

$$\tilde{T}^{\mu\nu} = \tilde{\rho} v^\mu v^\nu - \tilde{p}(g^{\mu\nu} - v^\mu v^\nu) + \tilde{\Pi}^{\mu\nu}, \quad (16)$$

where $\tilde{\Pi}^{\mu\nu}$ is the stress tensor and the four-velocity v^μ satisfies the condition

$$g_{\mu\nu} v^\mu v^\nu = 1. \quad (17)$$

In particular, the decomposition (16) holds true if the dark matter is determined by a phase space distribution Ω . In a homogeneous universe described by the metric (15) the stress tensor $\tilde{\Pi}^{\mu\nu}$ is absent. We introduce the energy momentum tensor T_Λ of an ideal fluid (with zero stress tensor) describing the dark energy

$$T_\Lambda^{\mu\nu} = (\rho_\Lambda + p_\Lambda) u^\mu u^\nu - g^{\mu\nu} p_\Lambda, \quad (18)$$

where $u^\mu u_\mu = 1$.

We consider a total energy momentum tensor which is conserved and consists of many components. We assume the energy momentum (non)conservation law (with certain currents N_b^ν ; the constant κ is describing a dissipation)

$$(T_b^{\mu\nu})_{;\mu} = 3\kappa^2 N_b^\nu. \quad (19)$$

We require

$$\sum_b N_b^\nu = 0, \quad (20)$$

if the total energy momentum is to be conserved.

In our diffusion model of sec.4 DM current is conserved

$$(N^\mu)_{;\mu} = 0 \quad (21)$$

This condition discriminates the model [3][4][28](introduced also in [29]) from other models of interacting DM/DE systems where N_b^ν are linearly related to the energy-momentum [30][31][32]. Eq.(21) in a homogeneous flat universe implies

$$N^0 = a^{-3} \frac{\tilde{\gamma}}{3\kappa^2}, \quad (22)$$

where we introduced another constant $\tilde{\gamma}$. N^0 has an interpretation of the particle density. Eq.(19) means that the increase of the energy density in an expanding universe is proportional to the particle density.

In order to satisfy the total energy-momentum conservation (20) we set

$$\gamma_\Lambda = -\tilde{\gamma} \equiv -\gamma. \quad (23)$$

Then, eqs.(19)-(23) in the frame $u = (1, \mathbf{0})$ and in a homogeneous space give

$$\partial_t \rho_\Lambda + 3H(1+w)\rho_\Lambda = -\gamma a^{-3}, \quad (24)$$

where $H = a^{-1}\partial_t a$ and

$$w = \frac{p_\Lambda}{\rho_\Lambda}.$$

The solution of eq.(24) for a constant w is

$$\rho_\Lambda(t) = a^{-3(1+w)}\sigma_\Lambda(0) - \gamma a(t)^{-3(1+w)} \int_{t_0}^t a(s)^{3w} ds \quad (25)$$

$\sigma_\Lambda(0)$ is a constant such that

$$\rho_\Lambda(t_0) = a(t_0)^{-3(1+w)}\sigma_\Lambda(0)$$

($\sigma_\Lambda(0)$ is just the initial DE density if we choose the convention $a(t_0) = 1$). In the frame $v = (1, \mathbf{0})$ eq.(19) for $\tilde{T}^{\mu 0}$ reads

$$\partial_t \tilde{\rho} + 3(1+\tilde{w})H\tilde{\rho} = \gamma a^{-3} \quad (26)$$

If we approximate $\tilde{w}$ as a constant then we get an analog of eq.(25)

$$\tilde{\rho}(t) = a^{-3(1+\tilde{w})}\tilde{\sigma}(0) + \gamma a(t)^{-3(1+\tilde{w})} \int_{t_0}^t a(s)^{3\tilde{w}} ds. \quad (27)$$

For the relativistic diffusion of sec.4 $\tilde{w}$ is time dependent $0 < \tilde{w}(a) \leq \frac{1}{3}$. As discussed in the next section $\tilde{w}(a) = \frac{1}{3}$ for massless particles. For $\tilde{w}(a) = \frac{1}{3}$

$$\tilde{\rho}(t) = a^{-4}\tilde{\sigma}(0) + \gamma a(t)^{-4} \int_{t_0}^t a(s) ds.$$

It is interesting to calculate the energy densities (27) for various expansion laws of $a(t)$. So, if $\tilde{w} > 0$ and $a(t) = t^\alpha$ then the diffusive term in eq.(27) (proportional to γ) is dominating the expansion for a large time as $\tilde{\rho} \simeq \gamma a^{-3+\frac{1}{\alpha}}$. This domination remains true for the exponential expansion, if $a = \exp(Ht)$, then $\tilde{\rho} \simeq \gamma a^{-3}$.

We apply the equilibrium thermodynamic definition of the entropy (due to Clausius and Caratheodory) as the integrating factor for the heat Q_b to each

component b of the cosmic fluid. Then, we have the Gibbs-Duhem differential (no chemical potential)

$$TdS_b = dQ_b = dE_b + p_b dV = (\rho_b + p_b)dV + V d\rho_b \quad (28)$$

with $p dV$ denoting the work done during the expansion of the volume $V = V_0 \sqrt{g}$ where $g = |\det(g_{\mu\nu})|$. Using

$$\partial_\mu \sqrt{g} = \Gamma_{\alpha\mu}^\alpha \sqrt{g}, \quad (29)$$

where Γ is the Christoffel symbol, from eqs.(26) and (28) in the frame moving with the fluid ($v = (1, 0)$) we obtain for an ideal fluid of DM

$$Td\tilde{S} = (\partial_\tau \tilde{\rho} + \Gamma_{\alpha 0}^\alpha (\tilde{\rho} + \tilde{p})) V_0 \sqrt{g} dt = 3\kappa^2 \tilde{N}^0 V_0 \sqrt{g} dt. \quad (30)$$

Hence, from eq.(22) in the homogeneous flat universe

$$Td\tilde{S} = V_0 \gamma dt. \quad (31)$$

Let us compare now the formulae for the energy (7) and (27). We can see that B in eq.(7) cannot be a constant if there is an energy non-conservation. In the derivation of ρ in sec.2 we should take into account that ρ depends on two variables a and A where

$$A = \int_{t_0}^t a^{3w} ds.$$

We can write $\tilde{\rho}$ (27) in the form

$$\tilde{\rho} = 3Z B(\tilde{A}) V_0^{-1} T_0^{-\frac{1+\tilde{w}}{w}} T_a^{\frac{1+\tilde{w}}{w}}, \quad (32)$$

where we denoted

$$Z = \frac{\tilde{\sigma}(0) V_0}{3T_0}$$

and $\kappa^2 = \frac{\gamma}{3Z}$ (we use this complicated way of expressing the energy density (27) in order to comply with the constants appearing in the diffusion of sec.4, Z is the statistical sum in the model of sec.4) T_a is the "temperature at fixed a " determined by eq.(9) and

$$B(\tilde{A}) = \frac{\tilde{\sigma}(0) V_0}{3Z} + \frac{\gamma V_0}{3Z} \int_{t_0}^t a(s)^{3\tilde{w}} ds \equiv T_0 + \kappa^2 \tilde{A} \quad (33)$$

Next, we must modify the formula for the entropy (8) derived through an "adiabatic" variation of variables. We assume that the entropy $\tilde{S}$ consists of the "adiabatic" part $\tilde{S}_{ad}$ (8) and the dissipative part $\tilde{S}_\gamma$ (31). Hence, taking into account a varying V as in eqs.(3) and (5) and integrating eq.(31) we obtain

$$\tilde{S}(t) = \tilde{S}_{ad} + \tilde{S}_\gamma = V(1 + \tilde{w}) \frac{\tilde{\rho}}{T} + V_0 \gamma \int_{t_0}^t \frac{ds}{T(s)} \quad (34)$$

where $V = V_0 a^3$.

If we demand that $\tilde{S}_{ad}$ does not change during the expansion then using the result (32) we obtain the dependence of T on a

$$T = B(\tilde{A}) T_0^{-1} T_a \quad (35)$$

An insertion of eq.(35) in eq.(34) gives

$$\tilde{S} = 3Z \ln B(\tilde{A}) + V(1 + \tilde{w}) \frac{\tilde{\rho}}{T} = 3Z \ln \left(\frac{T T_0}{T_a} \right) + V(1 + \tilde{w}) \frac{\tilde{\rho}}{T}. \quad (36)$$

Eq.(36) agrees with the definition of temperature [33]

$$\frac{1}{T} = \frac{1}{V} \frac{\partial \tilde{S}}{\partial \tilde{\rho}} \quad (37)$$

In sec.5 we calculate the entropy (here and in further discussions we set the Boltzmann constant $k_B = c = \hbar = 1$)

$$\tilde{S} = - \int d\mathbf{x} \frac{d\mathbf{p}}{(2\pi)^3} g \Omega \ln \Omega \quad (38)$$

for a system of diffusing particles with a probability distribution $\Omega(\mathbf{x}, \mathbf{p})$ in the phase space. We show that the formula (35) for temperature and the formula (32) for the energy density are satisfied for this realization of the dissipative system.

Writing an analog equation for $T_\Lambda dS_\Lambda$ we get eqs.(33)-(34) with $\gamma \rightarrow -\gamma$. The variation of the total entropy $\tilde{S} + S_\Lambda$ during universe expansion depends on the scale factor a and will be discussed in sec.6. In the generalized second law of thermodynamics [17][18][19][34][35] the entropy of the apparent horizon plays an important role contributing to the entropy increase [36] and compensating the entropy decrease during expansion of other components. However, in models of this section the entropy of the dark sector is increasing (see sec.6). So, the increase of the entropy of the horizon does not play such a decisive role. In the Appendix we generalize the derivations of this section to an arbitrary interaction N^ν in eq.(19) (without an assumption of the current conservation (21)). However, so far we have no candidate for N^ν other than the particle current.

4 Relativistic diffusion

In this section we discuss a Markovian approximation of an interaction of the system (DM) with an environment (DE) which leads to a description of this interaction by a diffusion. We consider a relativistic generalization of the Kramers diffusion defined on the phase space. It is determined in the unique way

by the requirement that the diffusing particle moves on the mass-shell $p^2 = m^2$ (see [37][38][39][40]).

In the contravariant spatial coordinates p^j on the mass shell we define the Riemannian metric G_{jk}

$$ds^2 = g_{\mu\nu} dp^\mu dp^\nu = -G_{jk} dp^j dp^k,$$

where p_0 is expressed by p^j .

The inverse matrix is (we assumed that $g_{0k} = 0$)

$$G^{jk} = -g^{jk} m^2 + p^j p^k. \quad (39)$$

Then,

$$\Gamma \equiv \det(G_{jk}) = \det(g_{jk}) p_0^{-2},$$

where

$$p_0^2 = m^2 - g_{jk} p^j p^k.$$

We define diffusion as a stochastic process generated by the Laplace-Beltrami operator Δ_H^m on the mass shell

$$\Delta_H^m = \frac{1}{\sqrt{\Gamma}} \partial_j G^{jk} \sqrt{\Gamma} \partial_k, \quad (40)$$

where $\partial_j = \frac{\partial}{\partial p^j}$.

The transport equation for the diffusion generated by Δ_H reads

$$(p^\mu \partial_\mu^x - \Gamma_{\mu\nu}^k p^\mu p^\nu \partial_k) \Omega = \kappa^2 \Delta_H^m \Omega, \quad (41)$$

where κ^2 is the diffusion constant and $\partial_\mu^x = \frac{\partial}{\partial x^\mu}$.

The relativistic diffusion determines the phase space distribution. Then, we can define the current

$$N^\mu = \sqrt{g} \int \frac{d\mathbf{p}}{(2\pi)^3} p_0^{-1} p^\mu \Omega, \quad (42)$$

where $g = |\det(g_{\mu\nu})|$ and the energy-momentum

$$\tilde{T}^{\mu\nu} = \sqrt{g} \int \frac{d\mathbf{p}}{(2\pi)^3} p_0^{-1} p^\mu p^\nu \Omega. \quad (43)$$

It can be shown for a general metric that the relation (19) is satisfied [28]

$$(\tilde{T}^{\mu\nu})_{;\mu} = 3\kappa^2 N^\nu, \quad (44)$$

where the current N^μ is conserved

$$(N^\mu)_{;\mu} = g^{-\frac{1}{2}} \partial_\mu (g^{\frac{1}{2}} N^\mu) = 0. \quad (45)$$

From eq.(45) we get eq.(22) in a homogeneous metric. If Ω is x -independent and $\int d\mathbf{x} = V_0$ then the constant γ can be expressed from eq.(42) (according to eq.(22)) as

$$\frac{\gamma}{3\kappa^2} = N^0 \sqrt{g} = g \int \frac{d\mathbf{p}}{(2\pi)^3} \Omega \equiv Z V_0^{-1}, \quad (46)$$

where $Z \equiv \int d\mathbf{x} g \int \frac{d\mathbf{p}}{(2\pi)^3} \Omega$ is the statistical sum.

In the massless case from $\tilde{T}_\mu^\mu = 0$ we obtain $\tilde{w} = \frac{1}{3}$, then from eq.(27)

$$\begin{aligned} \tilde{\rho}(t) &= a^{-4} \tilde{\sigma}(0) + \gamma a(t)^{-4} \int_{t_0}^t a(s) ds \equiv 3Z V_0^{-1} a^{-4} \left(\frac{V_0}{3Z} \tilde{\sigma}(0) + \kappa^2 \tilde{A} \right) \\ &= 3Z V^{-1} a^{-4} (T_0 + \kappa^2 \tilde{A}), \end{aligned} \quad (47)$$

where $T_0 \equiv \tilde{\sigma}(0) \kappa^2 \gamma^{-1} = V_0 \tilde{\sigma}(0) (3Z)^{-1}$.

We can solve eq.(41) exactly when $m = 0$ and the initial condition is the Jüttner distribution [41]. The solution is [3]

$$\Omega^J(t, T_0) = C_R T_0^3 (T_0 + \kappa^2 \tilde{A})^{-3} \exp \left(- \frac{a^2}{T_0 + \kappa^2 \tilde{A}} |\mathbf{p}| \right), \quad (48)$$

where T_0 is a free parameter such that $T_0 a(t_0)^{-1}$ can be interpreted as a temperature at $t = t_0$ (we assume $a(t_0) = 1$; from eq.(47) T_0 is related to $\tilde{\sigma}(0)$ and to Z). We introduced a constant C_R which determines the normalization of Ω^J . If we normalize Ω^J to the particle density n at t_0

$$\int d\mathbf{p} \frac{1}{(2\pi)^3} g \Omega^J(T_0) = n \equiv \frac{N}{V_0}, \quad (49)$$

where N is the number of particles, then

$$C_R = n \pi^2 \frac{1}{T_0^3}. \quad (50)$$

Summing over the temperatures T_0 with a density $q(T_0)$

$$\Omega(t) = \int dT_0 q(T_0) \Omega^J(T_0, t) \quad (51)$$

we get a more general solution of the diffusion equation.

In [40] we have derived an exact solution of the diffusion equation (41) with $m \geq 0$ in de Sitter space when $a(t) = \exp(H(t - t_0))$. Then, the solution is

$$\Omega(t) = C_R \exp(-3Ht) \exp \left(- \frac{H}{\kappa^2} \sqrt{a^2 \mathbf{p}^2 + m^2} \right). \quad (52)$$

In [42] we have shown that if $a(t)$ is close to an exponential function then we can obtain an expansion around the solution (52).

We can calculate now explicitly from eq.(48)

$$N^0 = C_R \frac{1}{(2\pi)^3} 8\pi T_0^3 a^{-3}. \quad (53)$$

From

$$\partial_\tau \tilde{T}^{00} + 4H\tilde{T}^{00} = \gamma a^{-3}$$

and eq.(47) we obtain

$$\gamma = \frac{1}{(2\pi)^3} 24\pi\kappa^2 T_0^3 = 3\kappa^2 Z_J V_0^{-1} = 3\kappa^2 n. \quad (54)$$

It is still interesting to consider the non-relativistic limit of eq.(41)[3][4]

$$m^2 a^2 \rightarrow \infty. \quad (55)$$

Then, the limit of eq.(41) is

$$m^{-1} \kappa^{-2} (\partial_\tau - 2H p^j \partial_j) \Omega = a^{-2} \Delta_{\mathbf{p}} \Omega, \quad (56)$$

where $\Delta_{\mathbf{p}}$ is the Laplacian in the momentum space. This is the non-relativistic diffusion equation in an expanding momentum space (see [43]). A special solution (with a Maxwell-Boltzmann initial condition) of the diffusion equation (56) reads

$$\Omega_{NR}(t, T_0, \mathbf{p}, \mathbf{k}) = C_{NR} T_0^{\frac{3}{2}} (T_0 + \kappa^2 A_{NR})^{-\frac{3}{2}} \exp\left(-\frac{(a^2 \mathbf{p} - a^2 \mathbf{k})^2}{2m(T_0 + \kappa^2 A_{NR})}\right), \quad (57)$$

where

$$C_{NR} = n T_0^{-\frac{3}{2}} (2\pi m)^{\frac{3}{2}}$$

which follows from the normalization

$$Z = \int \frac{d\mathbf{p}}{(2\pi)^3} g \Omega = n$$

and

$$A_{NR} = 2 \int_0^t a^2(s) ds. \quad (58)$$

The solution with a general initial condition is

$$\Omega_t(\mathbf{p}) = \int d\mathbf{k} dT_0 \Omega_{NR}(t, T_0, \mathbf{p}, \mathbf{k}) q(T_0, \mathbf{k}).$$

5 Entropy, free energy and temperature of the diffusive dark matter

We have discussed the first law of equilibrium thermodynamics in sec.2. The expansion of the universe can hardly be considered as a sequence of infinitesimal transitions between equilibrium states. For an expanding universe we need a formulation of non-equilibrium thermodynamics. If a phase space evolution Ω

is given then we can define the relative entropy for any two solutions Ω and Ω_J of the diffusion equation (41)[13][14]

$$\mathcal{F} = Z^{-1} \frac{1}{(2\pi)^3} \int d\mathbf{x} d\mathbf{p} g \Omega \ln \left(Z^{-1} Z_J \Omega \Omega_J^{-1} \right), \quad (59)$$

where

$$Z = \frac{1}{(2\pi)^3} \int d\mathbf{x} d\mathbf{p} g \Omega, \quad (60)$$

$$Z_J = \frac{1}{(2\pi)^3} \int d\mathbf{x} d\mathbf{p} g \Omega_J. \quad (61)$$

It can be shown that owing to the conservation of the current N^μ the normalizing integrals Z and Z_J do not depend on time. We have (the proofs follow ref.[14]; a slight extension is needed taking into account the time dependence of the metric)

$$\mathcal{F}(\Omega) \geq 0 \quad (62)$$

$\mathcal{F}(\Omega_J) = 0$ and

$$\partial_t \mathcal{F} = -Z^{-1} \frac{1}{(2\pi)^3} \int d\mathbf{x} d\mathbf{p} g \Omega G^{jk} \partial_j \ln Q \partial_k \ln Q \leq 0 \quad (63)$$

where

$$Q = \Omega \Omega_J^{-1}. \quad (64)$$

Let us consider in more detail the limit $a^2 m^2 \rightarrow 0$ in the diffusion equation (41). We choose the solution $\Omega^J(T_0)$ (48) as Ω^J in $\mathcal{F}$. Define

$$\Phi = Z\mathcal{F} - Z \ln(Z_J Z^{-1}) - 3Z \ln \left(C_R^{-\frac{1}{3}} T_0^{-1} (T_0 + \kappa^2 \tilde{A}) \right). \quad (65)$$

Then,

$$\partial_t \Phi \leq 0$$

Consider

$$U = \frac{a\tilde{E}}{(T_0 + \kappa^2 \tilde{A})} \equiv \frac{\tilde{E}}{T}, \quad (66)$$

where

$$\tilde{E} = \int d\mathbf{x} \sqrt{g} \tilde{T}^{00} = \frac{1}{(2\pi)^3} \int d\mathbf{p} d\mathbf{x} p^0 g \Omega = \int d\mathbf{x} \sqrt{g} \tilde{\rho}. \quad (67)$$

It follows from eq.(65) and (59) that $\tilde{F} = \Phi T$ satisfies the thermodynamic identity defining the free energy $\tilde{F}$

$$\frac{\tilde{F}}{T} = -\tilde{S} + U = -\tilde{S} + \frac{1}{T} \tilde{E}, \quad (68)$$

where $\tilde{S}$ is defined in eq.(38) and

$$T(t) = \frac{T(t_0) + \kappa^2 \tilde{A}}{a}. \quad (69)$$

The formula (69) for the temperature results from eqs.(37) and (68) through the basic principles of statistical thermodynamics [33] (it also follows from the Jüttner distribution (48))

$$\frac{1}{T(t)} = \frac{\partial \tilde{S}}{\partial \tilde{E}}. \quad (70)$$

For a powerlike evolution the temperature (69) is increasing and $T(t) \rightarrow \infty$ as $t \rightarrow \infty$. However, the powerlike evolution is considered in cosmology only for an intermediate time. For a large time we expect constant acceleration $a(t) = \exp(Ht)$ resulting from the cosmological constant. Then, the temperature tends to a finite value

$$T(t) \rightarrow \kappa^2 H^{-1}. \quad (71)$$

We could calculate this temperature inserting the formula (52) for Ω in de Sitter space into the formula (38) for the entropy and calculate T from eq.(70).

Let us calculate the entropy of Ω^J from eq.(38)

$$\tilde{S}^J = 3nV_0 - nV_0 \left(\ln(n\pi^2) - 3 \ln(T_0 + \kappa^2 A) \right). \quad (72)$$

This formula coincides (up to a constant) with eq.(34) derived on the basis of the first law of thermodynamics. The dependence of entropy on temperature (70) coincides with the standard one for Jüttner distribution (although now the temperature depends on time according to eq.(69)).

We can repeat the formulation of the non-equilibrium thermodynamics in the non-relativistic limit $a^2 m^2 \rightarrow \infty$ defining

$$\mathcal{F}(\Omega) = (2\pi)^{-3} Z^{-1} \int d\mathbf{x} d\mathbf{p} g \Omega \ln(Z^{-1} \Omega Z_{NR} \Omega_{NR}^{-1}) \quad (73)$$

where Ω_{NR} denotes the Maxwell-Boltzmann distribution (57) with $\mathbf{k} = 0$. We have $\mathcal{F}(\Omega_{NR}) = 0, \mathcal{F} \geq 0$ and $\partial_t \mathcal{F} \leq 0$. We define

$$\Phi = Z\mathcal{F} - Z \ln(Z_{NR} Z^{-1}) - \frac{3}{2} Z \ln \left(C_{NR} T_0^{-1} (T_0 + \kappa^2 A_{NR}) \right) \quad (74)$$

with

$$Z_{NR} = (2\pi)^{-3} \int d\mathbf{p} d\mathbf{x} g \Omega_{NR} = C_{NR} T_0^{\frac{3}{2}} (2\pi)^{-3} (2m\pi)^{\frac{3}{2}}$$

where

$$C_{NR} = n T_0^{-\frac{3}{2}} (2\pi)^{\frac{3}{2}} m^{\frac{3}{2}}$$

We have from the definition of $\mathcal{F}$ and the free energy $\tilde{F} = \Phi T$

$$\frac{\tilde{F}}{T} = -\tilde{S} + \frac{\tilde{E}_{nr}}{T} \quad (75)$$

where

$$\frac{\tilde{E}_{nr}}{T} = \frac{a^5 V_0}{T_0 + \kappa^2 A_{NR}} \tilde{\rho}_{nr} = \frac{V}{T} \tilde{\rho}_{nr} \quad (76)$$

and

$$\tilde{\rho}_{nr} = \sqrt{g} \int d\mathbf{p} \frac{\mathbf{p}^2 a^2}{2m} \Omega_{NR} \quad (77)$$

with the temperature defined by

$$\frac{1}{T} = \frac{\partial \tilde{S}}{\partial \tilde{E}_{nr}} = \frac{a^2}{T_0 + \kappa^2 A_{NR}}. \quad (78)$$

The a^{-2} decay of the temperature in the non-relativistic non-diffusive case $\kappa^2 = 0$ is well known [44]. The calculation of entropy from the Gibbs formula (38) gives for the Maxwell-Boltzmann temperature state (57) the result

$$S(\Omega_{NR}) = \frac{3}{2} Z_{NR} + \frac{3}{2} Z_{NR} \ln \left(C_{NR}^{-\frac{2}{3}} T_0^{-1} (T_0 + \kappa^2 A_{NR}) \right) + V \frac{\rho_{nr}}{T}, \quad (79)$$

which up to a constant agrees with the result (34) (with $\tilde{w} = 0$). Its dependence on temperature coincides with the standard formulas of classical statistical physics of ideal gases.

6 Evolution of the temperature and entropy

In this section we study the entropy and temperature evolution law in more detail. For arbitrary scale factor a the temperature (35) is an involved function of time. We have

$$T(t) = a^{-3\tilde{w}} \left(T_0 + \kappa^2 \int_{t_0}^t a^{3\tilde{w}}(s) ds \right) \quad (80)$$

and

$$\partial_t T = \kappa^2 - 3\tilde{w}HT.$$

For $\tilde{w} > 0$ (DM) and large t the temperature is an increasing function of time because $\int_{t_0}^t a^{3\tilde{w}} >> a^{3\tilde{w}}$ for $t >> t_0$.

From eq.(35) for DE ($\gamma \rightarrow -\gamma$) we obtain the formula

$$T_\Lambda = a^{3|w|} \left(T_{\Lambda 0} - \kappa^2 \int_{t_0}^t ds a^{3w} \right). \quad (81)$$

Hence,

$$\partial_t T_\Lambda = -\kappa^2 - 3wHT_\Lambda. \quad (82)$$

In general, the behaviour of the temperature depends on the form of the expansion factor. If we take $a = \exp(H(t - t_0))$ then, if $T_0 > T_\infty \equiv \frac{\kappa^2}{3Hw}$ then the

temperature of DM is monotonically decreasing to its minimal value $T_\infty = \frac{\kappa^2}{3\tilde{w}H}$. If $T_0 < T_\infty$ then the temperature of DM is monotonically increasing to its maximal value $\frac{\kappa^2}{3\tilde{w}H}$. For DE, if $T_{\Lambda 0} > \frac{\kappa^2}{3H|w|}$ then for the exponential expansion, the temperature is exponentially increasing to infinity. If $T_{\Lambda 0} < \frac{\kappa^2}{3H|w|}$ then the DE temperature becomes negative and the entropy does not make sense. For a powerlike expansion $a(t) = t_0^{-\alpha} t^\alpha$ the temperature of DM grows linearly. The temperature of DE is non-negative for all $t \geq t_0 > 0$ only if $3\alpha w + 1 < 0$ and $T_0 \geq \frac{\kappa^2}{|3w\alpha+1|} t_0$. Then, $\int_{t_0}^\infty ds a^{3w} < \infty$ in eq.(81). If $3\alpha w + 1 > 0$ then the temperature becomes negative for a large time and the entropy does not make sense.

The entropy of the DM/DE system is a sum of entropies. We do not know a phase space distribution of DE. However, from the energy conservation (eqs.(19)-(20) and (31)) it follows that

$$\partial_t S = \partial_t \tilde{S} + \partial_t S_\Lambda = V_0 \gamma \left(\frac{1}{T(s)} - \frac{1}{T_\Lambda(s)} \right). \quad (83)$$

Hence,

$$\partial_t S \geq 0$$

if $T \leq T_\Lambda$. This inequality is certainly satisfied for an exponential expansion (as long as the temperature makes sense) because we have the inequality

$$T_\Lambda \geq a^{3|w|} (T_{\Lambda 0} - \kappa^2 \int_{t_0}^\infty a^{3w}(s) ds) \quad (84)$$

and for an exponential expansion $T(t)$ is bounded. In general, we have $T(t) \leq Ct$ for a large time (with a certain constant $C > 0$). For a power law expansion we have additionally $C_0 t \leq T(t) \leq Ct$ with a certain $C_0 > 0$. Then, a sufficient condition for $T \leq T_\Lambda$ ensuring an increase of the entropy at large time is satisfied if

$$a^{3|w|} > Kt \quad (85)$$

with a sufficiently large K . For a powerlike expansion this means $3\alpha|w| > 1$; the condition which is also necessary for the temperature to be non-negative for all time. Note that the conditions $w < -\frac{1}{3}$ and $\alpha > 1$ are also needed for an accelerated expansion.

As in eq.(34) we can integrate eq.(83)

$$S = \tilde{S} + S_\Lambda = V_0 \gamma \int_0^t ds \left(\frac{1}{T(s)} - \frac{1}{T_\Lambda(s)} \right) + V(1 + \tilde{w}) \frac{\tilde{\rho}}{T} + V(1 + w) \frac{\rho_\Lambda}{T_\Lambda} \quad (86)$$

Using eqs.(80)-(81) (or returning to eq.(35)) we obtain

$$S = 3Z \ln \left(T_0 + \kappa^2 \int_{t_0}^t ds a(s)^{3\tilde{w}} \right) + 3Z \ln \left(T_{\Lambda 0} - \kappa^2 \int_{t_0}^t ds a(s)^{3w} \right) + V(1 + \tilde{w}) \frac{\tilde{\rho}}{T} + V(1 + w) \frac{\rho_\Lambda}{T_\Lambda}. \quad (87)$$

The first (logarithmic) term is increasing as a function of time whereas the second term is decreasing. If

$$\int_{t_0}^{\infty} ds a^{3w} < \infty \quad (88)$$

then the second term has a lower bound $\ln \left(T_{\Lambda 0} - \kappa^2 \int_0^{\infty} ds a(s)^{3w} \right)$. The condition (88) is necessary for a definition of entropy of DE as the argument of the logarithm must be positive. As discussed below eq.(82) the argument of the logarithm is negative for negative temperatures as can happen for DE with a powerlike expansion such that $3\alpha w > -1$. For systems which have a meaningful entropy of the dark sector the entropy increase is contained in the first term of eq.(87). The terms depending on the energy density in eq.(87) are constant as follows from the derivation of the temperature dependence in sec.2. It follows from eq.(83) that $\partial_t S > 0$ for $T < T_{\Lambda}$. As discussed in [36] the total entropy should tend to a maximum what requires $\partial_t^2 S_{tot} < 0$ where S_{tot} in [36] includes the entropy of the apparent horizon which has a negative second order derivative for some range of w_{eff} (where w_{eff} is the effective equation of state). In the case of the dark sector (86) we have from eqs.(80)-(82)

$$\partial_t^2 S = -\gamma \kappa^2 (T^{-2} + T_{\Lambda}^{-2}) + 3\gamma \tilde{w} \frac{H(t)}{T} - 3\gamma w \frac{H(t)}{T_{\Lambda}}. \quad (89)$$

The first term is negative whereas the remaining terms are positive. We need detailed calculations to estimate the balance of these terms. We have calculated these expressions for the exponential expansion $a = \exp(H(t - t_0))$ then

$$\partial_t^2 \tilde{S} = 3\tilde{w} H T^{-2} V_0 \gamma (T_0 - \frac{\kappa^2}{3\tilde{w}H}) \exp(-3\tilde{w}H(t - t_0)) \quad (90)$$

and

$$\partial_t^2 S_{\Lambda} = 3|w| H T_{\Lambda}^{-2} V_0 \gamma (T_0 - \frac{\kappa^2}{3|w|H}) \exp(3|w|H(t - t_0)). \quad (91)$$

For a large time

$$\partial_t^2 \tilde{S} \simeq (3\tilde{w}H)^3 \kappa^{-4} V_0 \gamma (T_0 - \frac{\kappa^2}{3\tilde{w}H}) \exp(-3\tilde{w}H(t - t_0)) \quad (92)$$

and

$$\partial_t^2 S_{\Lambda} \simeq 3|w| H V_0 \gamma \left(T_0 - \frac{\kappa^2}{3|w|H} \right)^{-1} \exp(-3|w|H(t - t_0)). \quad (93)$$

$\partial_t^2 S_{\Lambda} > 0$ because the dominator in eq.(93) must be positive if the temperature is to be positive. Then, the second time derivative of the entropy S can be negative only if $|w| > \tilde{w}$, so that the second derivative of S_{Λ} is decaying faster than the one of $\tilde{S}$. Moreover, for $\partial_t^2 S < 0$ we need

$$T_0 < T_{\infty} = \frac{\kappa^2}{3\tilde{w}H} \quad (94)$$

(then the temperature T is increasing to its maximal value T_∞). For power law expansion $a(t) = t_0^{-\alpha} t^\alpha$ we can show that (under the condition $3|w|\alpha > 1$ of non-negative temperature) $\partial_t^2 S_\Lambda$ is decaying faster than $\partial_t^2 \tilde{S}$. Hence, the latter is dominating at large time with the result

$$\partial_t^2 S \simeq -\gamma V_0 \kappa^{-2} (3\tilde{w} + 1) t^{-2} < 0. \quad (95)$$

The entropy slowly achieves its maximum at an infinite time.

7 Summary

It is known that negative pressure leads to a curious behaviour of thermodynamic functions of ideal non-interacting fluids. If we extend the argument to an expanding universe then we obtain a non-conventional dependence of temperature on the expansion scale factor a . In particular, the temperature of DE is increasing as a function of a . In this paper we assumed that the cosmological fluid consists of components which can exchange the energy. We have suggested a particular form of the energy (non) conservation assuming that the energy gain (or loss) is proportional to the particle density. We have obtained this energy exchange in a model of a relativistic diffusion of DM particles in a DE environment. We have shown that as a result of DM/DE interaction the dependence of the temperature on the scale factor undergoes a substantial change. The temperature of DM (which has positive pressure, $\tilde{w} > 0$) is growing to infinity as a function of time for power law expansion. For an exponential expansion it is reaching a fixed value $\frac{\kappa^2}{3\tilde{w}H}$ as time tends to infinity. The temperature of DE is increasing in all cases when the definition of temperature makes sense for arbitrarily large time. The entropy is increasing for a large time because the temperature of DM is increasing at a slower rate than the temperature of DE. This is a consequence of the energy transfer from DE to DM. We derive formulae for temperature and entropy in the case of more general DM/DE interactions. However, we do not have realizations of other interactions besides the one of the relativistic diffusion. The study of temperature, entropy and the heat flow in various components of the cosmological fluid should give important information about structure formation and its evolution.

Acknowledgement

The research is supported by the NCN grant DEC-2013/09/BST2/03455

8 Appendix

In the appendix we consider a more general system of DM/DE interactions. Let us denote $3\kappa^2 N^0 = \gamma Q$ in eq.(19). Then, in eq.(27)

$$\tilde{\rho}(t) = a^{-3(1+\tilde{w})} \tilde{\sigma}(0) + \gamma a(t)^{-3(1+\tilde{w})} \int_{t_0}^t a(s)^{3(1+\tilde{w})} Q(s) ds. \quad (96)$$

We write $\tilde{\rho}$ again in the form (32) with

$$B(\tilde{A}) = \frac{\tilde{\sigma}(0)V_0}{3Z} + \frac{\gamma V_0}{3Z} \int_{t_0}^t a(s)^{3(1+\tilde{w})} Q(s) ds \equiv T_0 + \kappa^2 \tilde{A}, \quad (97)$$

where $\kappa^2 = \frac{\gamma V_0}{3Z}$ and

$$\tilde{A} = \int_{t_0}^t a(s)^{3(1+\tilde{w})} Q(s) ds.$$

Demanding that S_{ad} in eq.(34) does not depend on time and using eq.(96) we derive the evolution of temperature

$$T = a(s)^{-3\tilde{w}} \left(T_0 + \frac{\gamma V_0}{3Z} \int_{t_0}^t a(s)^{3(1+\tilde{w})} Q(s) ds \right). \quad (98)$$

The formulae for DE are obtained from eqs.(96)-(98) by $\gamma \rightarrow -\gamma$ and $\tilde{w} \rightarrow w$. The entropy is

$$S = \tilde{S} + S_\Lambda = 3Z \ln \left(T_0 + \kappa^2 \tilde{A} \right) + 3Z \ln \left(T_{\Lambda 0} - \kappa^2 A \right) + V(1 + \tilde{w}) \frac{\tilde{\rho}}{T} + V(1 + w) \frac{\rho_\Lambda}{T_\Lambda} \quad (99)$$

where

$$A = \int_{t_0}^t a(s)^{3(1+w)} Q(s) ds,$$

The last two terms in eq.(99) are time independent. As can be seen from sec.6 the conclusions concerning the growth of temperature and entropy depend on the sign of γ and on whether Q grows or decays as a function of a .

References

- [1] P.A.R. Ade, et al arXiv:1403.3985[astro-ph.CO]
- [2] R.C.Tolman, Relativity, Thermodynamics and Cosmology, Oxford University Press, 1934
- [3] Z. Haba, Class.Quant.Grav.**31**,075011(2014)
- [4] Z.Haba, A. Stachowski and M. Szydlowski,JCAP07(2016)024
- [5] M. Planck, Vorlesungen Über Die Theorie Der Wärmestrahlung,J.A. Barth Verlag, Leipzig,1906
- [6] S. Weinberg, Astroph. J.**168**,175(1971)
- [7] S. Frautschi, Science **217**,593(1982)

- [8] P.T.Landsberg and D. Park, Proc.R.Soc.Lond.**A346**,485(1975)
- [9] O. Gron, Entropy **14**,2456(2012)
- [10] D. Wallace, Br.J.Phil.Soc.**61**,513(2010)
- [11] C.A. Egan and C.H. Lineweaver, Astroph.J.**710**,1825(2010)
- [12] W.Fischler and L. Susskind, arxiv:9806039
- [13] J.L. Lebowitz and P.G. Bergmann,Ann.Phys.(N.Y.)**1**,1(1957)
- [14] H. Risken, The Fokker-Planck Equation, Springer, 1989
- [15] B. Basu and D. Lynden-Bell, Q.J.R.Astr.Soc.**31**,359(1990)
- [16] A. Faltenbacher, Y. Hoffman, S. Gottlöber and G. Yepes, MNRAS**376**,1327(2007)
- [17] J. Bekenstein, Phys.Rev.**D9**,3292(1974),ibid **D12**,3077(1975)
- [18] G. Gibbons and S.W. Hawking, Phys.Rev. **D15**,2738(1977)
- [19] P.C. Davis and T.M. Davis, Found.Phys.**32**,1877(2003)
- [20] R. Maartens, arXiv:9609119
- [21] M.O. Calvao, J.A.S. Lima and I.Waga, Phys.Lett.**A162**,223(1992)
- [22] R.Silva, J.A.S. Lima and M.O. Calvao, Gen.Rel.Grav.**34**,865(2002)
- [23] H.H.B. Silva, R. Silva, R.S. Gonsalves, Zhong-Hong Zhu and J.S. Alcaniz, Phys.Rev.**D88**,127302(2013)
- [24] Yu.L. Bolotin,A. Kostenko,O.A. Lemets and D.A. Yerokhin, Int.J.Mod.Phys.**D24**,1530007(2014)
- [25] J.A.S. Lima,Phys.Rev.**D54**,2571(1996)
- [26] S.H. Pereira and J.F. Jesus,Phys.Rev.**D79**,043517(2009)
- [27] D. Pavon and Bin Wang, Gen.Relativ.Grav.**41**,1(2009)
- [28] Z.Haba,Mod.Phys.Lett.**A25**,2681(2010)
- [29] S. Calogero, J.Cosmo.Astrop.Phys.**11**,025(2011)
- [30] L. Amendola,arXiv 0610806
- [31] Ch.G. Böhmer, G. Caldera-Cabral, R. Lazkoz and R. Maartens, Phys.Rev.**D78**,023505(2008)

- [32] J.A.S. Lima and I. Baranov, Phys.Rev.**D90**,043515(2014)
- [33] L.Landau and E.M. Lifszic, Statistical Physics, Pergamon Press, Oxford,1980
- [34] R.G. Cai and S.P. Kim, JHEP02(2005)050
- [35] K.Karami, S. Ghaffari and M.M. Soltanzadeh,
Class.Quant.Grav.**27**,205021(2010)
- [36] N. Radicella and D. Pavon, Gen.Relativ.Gravit.**44**,685(2012)
- [37] R.Dudley, Arkiv for Matematik,**6**,241(1965)
- [38] J. Franchi and Y.Le Jan, Comm.Pure Appl.Math.**60**,187(2007)
- [39] Z. Haba, Phys.Rev.**E79**,021128(2009)
- [40] Z. Haba, Class.Quant.Grav.**27**,095021(2010)
- [41] F. Jüttner, Ann.Phys.(Leipzig)**34**,856(1911)
- [42] Z.Haba, Mod.Phys.Lett.**A31**,1650146(2016)
- [43] S.B. Yuste,E. Abad and C. Escudero, Phys.Rev**E94**,032118(2016)
- [44] J. Bernstein, Kinetic Theory in an Expanding Universe, Cambridge,1988